

The Consistency of Sex Ratio of *Drosophila melanogaster* (Meigen) in Different Physical Environment Condition

Ahmad Fauzi***, Shefa Dwijayanti Ramadani***, Ika Sukmawati ****

*Al Ya'lu Middle School

** Departement of Biology Education, Universitas Muhammadiyah Malang

***Departement of Biology Education, Universitas Islam Madura

**** Departement of Biology Education, Universitas Negeri Malang

Article Info

Article history:

Received Jul 12th, 2017

Revised Aug 20th, 2017

Accepted Oct 26th, 2017

Keyword:

Drosophila melanogaster

EMF exposure

Dark condition

High temperature

Sex ratio

ABSTRACT

Several environmental factors have been reported affect sex ratio in many organisms. The environmental factors that affecting sex ratio could from physical, chemical, or biological factors. In this study, the effect of some physical factors on sex ratio in *D. melanogaster* were studied. In normal laboratory condition, high temperature, dark condition, and EMF exposure, the sex ratio were 0.91:1.09, 1.00:1.00, 0.89:1.11, and 1.01:0.99 respectively. From the result, it can be seen that males and females adult filial number ratios in each group were close to 1:1.

Copyright © 2017 Green Technology.
All rights reserved.

Corresponding Author:

Ahmad Fauzi,

Al Ya'lu Middle School

Jl. Teluk Mandar No. 55 Arjosari, Malang, East Java, indonesia, 65126

Email: fauizou91@gmail.com

1. INTRODUCTION

Sex ratio is a topic often discussed in various biological studies. Sex ratio itself is a term that generally refers to the ratio of males to females in a population [1,2]. The ratio is commonly expressed as the number of males per 100 or 1000 females or the fraction of the total population that is female (or male) [1,3,5]. The proportion of females in the ratio indicate the reproductive potential of a population [4]. Moreover, sex ratio also reflects a number of life history traits, such as the importance of sexual mating system and also the other information related to the past, present, and future of a population [1,4].

In many organism with the sexes are separate, females are produced in approximately equal numbers with males [6]. Therefore, in many organism, such as insects, the sex ratio is 1:1 [5]. This ratio maximizes the availability of males to females, and hence maximizes genetic heterogeneity [4]. In the absence of manipulation, the sex ratio are constant [7]. However, several environmental factors ultimately affect sex ratios [4,6,8]. The environmental factors affecting sex ratio can from physical, chemical, or biological factors [1,4,6,8,9].

In this study, the effect of some physical factor on sex ration in *D. melanogaster* were studied. Those physical factor are environmental temperature, minimum light condition, and electromagnetic field (EMF) radiation. Temperature is one of the major factor regulating population growth [10,11]. This factor is one of the most important parameters affecting the live of organisms [12]. Beside temperature, light can also be a limiting factor [10,13]. Light can affect the reproductive of organisms [14]. On the other hand, recently, EMF field radiation have also been reported have the poteintal in affecting the population of organisms [15].

D. melanogaster was used as a model organism in this study. The organism was choosen in this study because of some reasons. First, *Drosophila* is an organism that produced consistently high numbers of offspring [16]. Second, this organism is recorded as rapid breeder with a lot of eggs and short life cycle [16-18]. Third, this organism has often been as model organism in many study examining various problems in biology [16,17].

2. RESEARCH METHOD

2.1. The organsim and environmental conditions

D. melanogaster wild-type strain from Genetic Laboratory FMIPA UM were used in this study. Flies were cultured in a 200 ml cylindrical glass bottle, with 7 cm diameter and 9 cm height, filled with 30 ml standard food, as described in Fauzi et al [15]. The flies cultures were kept in a research room at Genetic Laboratory FMIPA UM. When there were blackened pupae, the pupae were isolated into plastic tube with 1 cm diameter and 5 cm height. Adult flies that emerge from this plastic tube were using for crossbreeding at treatment stage.

2.2. Treatment stage

D. melanogaster ♂ wild-type strain was crossed with ♀ wild-type strain in glass bottle that the same with cultured bottle, both of it 2 X 24 hours after hatching from pupae. For temperature treatment, a pair of three replicates of *D. melanogaster* crossbred were exposed continuously to a constant temperature of 30° C in an incubator. For minimum light treatment, a pair of three replicates of *D. melanogaster* crossbred were placed in a sealed opaque box. The Crossbreeding box was then coated with black plastic and placed in a room that was not much exposed to light. For EMF exposure treatment, a pair of three replicates of *D. melanogaster* crossbred were placed close to GSM mobile phone with provider that using frequencies at 900/1800 MHz (3G) and HSDPA 2100 (4G) as a source of exposure source. Along with treatment groups, the normal laboratory room condition group was also set up. The group was placed in room with temperatures fluctuating naturally following the condition of the surrounding temperature (25-30° C), wiht normal light conditions (light-dark change/LD 12:12h), and was not exposed by mobile phone.

2.3. Data collection

The number of adult filial males and females at each group were recorded. Then, the data were calculated to get ratio of males to females.

3. RESULTS AND ANALYSIS

Fig. 1 is the graphs show the comparison between male and female adult filial number of *D. melanogaster* whereas Table 1. show the sex ratio at each different environmental condition. Based on Fig. 1 and Table 1, it can be seen that the comparison between males and females in each group were close to 1:1. The present results indicate sex ratio in *D. melanogaster* was stable on various environmental condition that have been designed in this study.




Figure 1. The comparison between male and female adult filial number of *D. melanogaster* at different environmental condition

The determination of sex in *Drosophila* is not only based on the existence of Y chromosome in it body. [19]. The determination of sex in this organism is more determined by the ratio of X chromosome number with the number of sets of autosomes [20]. This is due to the fact that the X chromosomes and autosomes together play a critical role in sex determination in *Drosophila* [19]. However, the determination of sex like that make the sex ratio *Drosophila* in normal condition still close to 1:1. But, the sex that determined by its genetic makeup or chromosomes in their body more refers to sex at embryo. After birth, or indeed before that, flies will growth and

develop until flies enter the adult phase, during the process some factor from environment will affect and control the growth of the population of the organism [10,11].

Table 1. Sex Ratio of *D. melanogaster* at Each Different Environmental Condition

| Group | Sex Ratio |
|-----------------------------|-----------|
| Normal laboratory condition | 0.91:1.09 |
| High temperature | 1.00:1.00 |
| Dark condition | 0.89:1.11 |
| EMF exposure | 1.01:0.99 |

Temperature is one of environmental factor that can influence sex determination in adult phase [19,20]. The temperatures can affect the survival level of males or females from larval phases until imago phases in such a way as to shift the sex ratio [19]. Furthermore, it has been known there are several gene in *Drosophila* that involve determining sex could be affected by temperature [21]. However, based on Fig. 1 and Table 1. in high temperature treatment, the sex ratio was still 1:1. The present result was also not agreement with previous researchs that reported temperature is one of environmental factor that can affect sex ratio in *Drosophila* [22]. But, the present result was in line with the other previous research reported there were no statistical support for an effect of temperature on the number of males and females reaching the adult stage [23]. Therefore, because of the sex ratio in *Drosophila* at embryo stage is 1:1 and there were no gender related effect of temperature during the organism growth until adult phase, the sex ratio in adult phase still 1:1.

In dark condition treatment, the sex ratio was 0.89:1.11. The ratio was still close to 1:1, but in this study, this ratio is the most deviates from 1:1. Until now, no reports have reported any effect of light on the sex ratio. Differences in the number of males and females in dark condition treatment in this study indicate there are gender related effect of light during the organism reach adult phase. Light is used by organisms as the external signal controlling circadian rhythms that regulating gene expression and physiological processes in their body [24-26]. Therefore, the differences in the number of males and females indicate a different degree of influence of light on circadian rhythms in both sexes.

In EMF exposure treatment, the sex ratio was 1.01:0.99. The ratio was very close to 1:1. This result was an agreement with previous study reported EMF may not exert a gender-related influence on flies [27]. Related to that, Moorhouse & Macdonald stated sex ratio variation is commonly observed as a consequence of varying survival and value of offspring of different sexes [28]. From the present result and the statement, it can be seen that, the radiosensitivity of male flies to EMF has the same level with female flies [27].

4. CONCLUSION

The effect of some physical factor on sex ration in *D. melanogaster* were conducted in the present study. In normal laboratory condition, high temperature, dark condition, and EMF exposure, the sex ratio were 0.91:1.09, 1.00:1.00, 0.89:1.11, and 1.01:0.99 respectively. From the result, it can be seen that the comparison between males and females in each group were close to 1:1. The results indicate sex ratio in *D. melanogaster* was stable on various environmental condition that have been designed in this study.

REFERENCES

- [1] Skalski RS, Ryding KE, Millspaugh JJ. Wildlife Demography: Analysis of Sex, Age, and Count Data. Burlington: Elsevier Academic Press; 2005. 49 p.
- [2] Bailey J. The Facts on File Dictionary of Ecology and the Environment. New York: Facts on File, Inc.; 2004. 214 p.
- [3] Qazi SA, Qazi NS. Population Geography. New Delhi: A. P. H. Publishing Corporation; 2006. 95 p.
- [4] Schowalter TD. Insect Ecology: An Ecosystem Approach, Fourth Edition. London: Academic Press; 2016. 147 p.
- [5] Prakahs M. Insect Behaviour. New Delhi: Discovery Publishing House Pvt. Ltd.; 2008. 245 p.
- [6] Hardy ICW, editor. Sex Ratios: Concepts and Research Methods. Cambridge: Cambridge University Press; 2002. 2 p.
- [7] Cherian A, Rose A, Gupta D, Minz S, Prasad J, George K. Whither Sex Ratios in a Low Mortality Setting. IJWHR [Internet]. 2016 April [cited 2017 June 28];4(2): 64-67. Available from: http://www.ijwhr.net/pdf/pdf_IJWHR_156.pdf DOI: 10.15296/ijwhr.2016.15

- [8] Rosenfeld CS, Roberts RM. Maternal Diet and Other Factors Affecting Offspring Sex Ratio: A Review. *Biology of Reproduction* [Internet]. 2004 June [cited 2017 June 28];71(4): 1063-1070. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/15229140> DOI: 10.1095/biolreprod.104.030890
- [9] Wajnberg E, Bernstein C, Alphen JV. *Behavioural Ecology of Insect Parasitoids: From theoretical approaches to Field Applications*. Malden: Blackwell Publishing; 2008. 254 p.
- [10] Miller GT, Spoolman SE. *Essentials of Ecology*. Belmont: Brooks/Cole Cengage Learning; 2009. 224 p.
- [11] Price MF, Byers AC, Friend DA, Kohler T, Price LW, editors. *Mountain Geography: Physical and Human Dimensions*. Berkeley: University of California Press; 2013. 224 p.
- [12] Bhatnagar VS. *Chandigarh, the City Beautiful: Environmental Profile of a Modern Indian City*. New Delhi: A. P. H. Publishing Corporation; 1996. 61 p.
- [13] Roberts M, Reiss M, Monger G. *Advanced Biology*. Chetenham: Nelson; 2000. 199.
- [14] Inyang HI, Daniels JL, editors. *Environmental Monitoring, Volume 1*. Oxford: Eolss Publishers Co. Ltd.; 2009. 22 p.
- [15] Fauzi A, Corebima AD, Zubaidah S. The Fluctuation of Adult Filial Number and Eclosion Time of *Drosophila melanogaster* that Exposed by Mobile Phone in Multiple Generations. In: the 6th Annual Basic Science International Conference [Internet]; 2016 March 2-3; Atria Hotel and Conference, Malang, Indonesia; Faculty of Mathematics & Sciences, Brawijaya University; 2016 June [cited 2017 June 28]; p: 124-128. Available from: <http://basic.ub.ac.id/files/proceeding/PROCEEDINGS-BASIC-2016.pdf>
- [16] Neethu BK, Babu YR, Harini BP. Flavors supplemented in diet regulate the hatchability and viability in *Drosophila*. *Dros. Inf. Serv* [Internet]. 2014 [cited 2017 June 28];2013(2013):24-28. Available from: <http://www.ou.edu/journals/dis/DIS97/Neethu%2024.pdf>
- [17] Jennings BH. *Drosophila – a versatile model in biology & medicine*. *Materialstoday* [Internet]. 2011 May [cited 2017 June 28]; 14(5): 190-195. Available from: <http://www.sciencedirect.com/science/article/pii/S1369702111701134> DOI: 10.1016/S1369-7021(11)70113-4.
- [18] Fauzi A, Corebima AD, Zubaidah S. The Utilization of *Drosophila melanogaster* as a Model Organism in Genetics I and Genetics II Courses in Faculty of Mathematics and Natural Science, State University of Malang. In: *International Conference on Education and Training 2016* [Internet]; 2016 Nov 4-6; Malang; Faculty of Education, State University of Malang; 2016 [cited 2017 June 28]; p: 51-56. Available from: <http://icet.fipum.com/wp-content/uploads/2017/03/PROCEEDINGS-2ND-ICET-BOOK-1-.pdf>
- [19] Precht H, Christophersen J, Hensel H, Larcher W. *Temperature and Life*. Berlin: Springer-Verlag; 1973. 382 p.
- [20] Egna HS, Boyd CE. *Dynamics of Pond Aquaculture*. Boca Raton: CRC Press; 1997. 234 p.
- [21] Feldmeyer BV. The effect of temperature on sex determination [thesis on internet]. Groningen: University of Groningen; 2009. [cited 2017 June 29]. Available from: <http://www.rug.nl/research/portal>
- [22] Battle AJX. *Fruit Fly Frenzie: The Effect of Temperature on Gender Development of Drosophila melanogaster* [Internet]. California State Science Fair, 2014 [cited 2017 June 29]. Available from: cssf.usc.edu/History/2014/Projects/J2202.pdf
- [23] Choi J, Han D, Yang W. The effect of temperature on the number and gender of fruit flies (*Drosophila melanogaster*) reaching adulthood. *The Expedition* [Internet]. 2014 [cited 2017 June 29];4(2014). Available from: <http://ojs.library.ubc.ca/index.php/expedition/article/download/186418/185443>
- [24] Zee PC, editor. *Science of Circadian Rhythms, An Issue of Sleep Medicine Clinics*. Philadelphia: Elsevier; 2015. 427.
- [25] Young MW, editor. *Methods in Enzymology: Circadian Rhythms*. San Diego: Elsevier Academic Press; 2005. 27.
- [26] Squire LR, editor. *Encyclopedia of Neuroscience: Volume One, Volume 1*. Oxford: Academic Press, 965 p.
- [27] Fauzi A. The Comparison of Males and Females Number of *Drosophila melanogaster* that Exposed by Mobile Phone in Multiple Generations. Paper presented at: 3rd IBOC 2016 & 10th Korea Asean Biomass Symposium; 2016 Oct 15; ITS Campus, Surabaya, Indonesia.
- [28] Moorhouse TP, MacDonald DW. Indirect negative impacts of radio-collaring: sex ratio variation in water voles. *Journal of Applied Ecology* [Internet]. 2005 Feb [cited 2017 June 29]; 42(1): 91-98. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2664.2005.00998.x> DOI: 10.1111/j.1365-2664.2005.00998.x