

Bacillus thuringiensis As Local Biological Agent

Akhmad Rizali¹

¹Department of Agrotechnology, Lambung Mangkurat University, Indonesia

Abstract- The chemical insecticides are still contributing to human life enormously, but they have been distributed in ecological system of organisms including human beings because of their low specific toxicity to any organism and their low specific toxicity to any organism and their slight decomposition in nature. Therefore, many biological control of insects have been investigated. Currently, researches on the use pathogenic microorganisms to control insect pests are increasing. Microbial pest control is practiced in different parts of the world though utilization of pathogen like fungi, bacteria, viruses and nematodes. Bacterial research causing disease in insects began in the late nineteenth century. It was a study of flacherie of the silkworm, *bombyx mori* in this report on the discovery of *sotto bacillus*, referred briefly to occurrence of sotto bacillus-like organism, which causes the disease to silkworm larvae. Ten-fold serial dilutions of the heated suspension in sterile distilled water were placed on nutrient agar (NA-pH 7.5). After two days of incubation at 28°C, *Bacillus* colonies were recorded. After 2 to 3 days incubation, crystalliferous sporeforming bacteria were determined in phase contrast microscop. Isolation from five soil samples yielded about 35 isolates, only one was identified as *B. thuringiensis*.

Keywords—*B. thuringiensis*. Biological Agent

I. INTRODUCTION

Cabbage moth, *C. binotalis* Zell. (Lepidoptera: Pyraustidae) is considered the most important limiting factor for a successful production of cruciferous vegetable not only in the Indonesia but in other country in the world. The larva feeds on foliage from seedling to harvest causing 100% yield loss if not control (Rejesus and Sayaboc, 1990).

Numerous chemical insecticides have been used in order to control pests, which damage for agriculture. While chemical insecticides have knock down effect to the insect pests, they are too expensive in the developing countries and harmful to both human and the environment. In addition, target insect pests rapidly develop biological resistance especially at higher rates of application. The chemical insecticides are still contributing to human life enormously, but they have been distributed in ecological system of organisms including human beings because of their low specific toxicity to any organism and their low specific toxicity to any organism and their slight decomposition in nature (Shorey and Hall, 1962). Therefore, many biological control of insects have been investigated. Currently, researches on the use pathogenic

microorganisms to control insect pests are increasing. Microbial pest control is practiced in different parts of the world though utilization of pathogen like fungi, bacteria, viruses and nematodes. Bacterial research causing disease in insects began in the late nineteenth century. It was a study of flacherie of the silkworm, *bombyx mori* (Burgess and Hussey, 1971; Burgess, 1981). Ishiwata (1901) in this report on the discovery of *sotto bacillus*, referred briefly to occurrence of sotto bacillus-like organism, which causes the disease to silkworm larvae.

Berliner (1911) proposed the name of *B. thuringiensis* for a species of bacillus which was isolated from the diseased larvae of the Mediterranean flour moth *Anagasta (Ephestia) kuehniella* Zell. Later, Berliner (1915) noted infection of the larvae after the ingestion of the bacillus or its spore, described and named it *Bacillus thuringiensis*. Mattes (1927) isolated the same bacillus from the same insect host, which Berliner had found earlier.

B. thuringiensis is a gram-positive soil bacterium, and produce a crystalline inclusion body during sporulation (Bulla et al., 1980). This parasporal body is composed of proteins termed “delta-endotoxin”, and specifically toxic to insects. In addition, *B. thuringiensis* produce another toxins namely: alpha-toxin, beta-exotoxin, and gamma-exotoxin. All of the toxic substance may not present in the bacterium (Heimpel, 1967). In another hand, Krieg (1961) has defined various toxic substance produced *B. thuringiensis* as follow: (a) thermolabile endotoxin; (b) thermostable exotoxin; (c) bacillogenic antibiotic; (d) lecithinase; (e) proteinase.

Most strains of *B. thuringiensis* produce delta-endotoxin crystals toxic to lepidopteran insects such as moth (Dulmage et al., 1970). Recently, however several researches have shown that *B. thuringiensis* is also widely distributed in natural soils of various area. Delucca et al., (1982) reported that *B. thuringiensis* made up less than 0.5% of more than 46,000 bacterial isolates recovered from various soils in the United States.

The objective of the studies to survey, collect and determine the distribution of *B. thuringiensis* in selected diverse crop-growing area.

II. MATERIAL AND METHOD

Isolation of *B. thuringiensis*